

Donahaye, E.J., Navarro, S. and Leesch J.G. [Eds.] (2001) *Proc. Int. Conf. Controlled Atmosphere and Fumigation in Stored Products, Fresno, CA. 29 Oct. - 3 Nov. 2000, Executive Printing Services, Clovis, CA, U.S.A pp. 231-239*

STORAGE OF DATES UNDER CARBON DIOXIDE ATMOSPHERE FOR QUALITY PRESERVATION

S. NAVARRO*, J.E. DONAHAYE, MIRIAM RINDNER AND A. AZRIELI

*Dept. of Stored Products, Agricultural Research Organization, The Volcani Center, Bet Dagan 50250, Israel [*e-mail: vtshlo@netvision.net.il]*

ABSTRACT

Dates grown in Israel served as a model to demonstrate the feasibility of storing dried fruit under CO₂ atmosphere. A controlled atmosphere of 60-80% CO₂ was used within a 151 m³ plastic chamber partially filled with 30 tonnes of dates stacked in crates on pallets. The dates were stored in the chamber in bulk (in boxes of 10 kg or crates of 400 kg on pallets) or packed (250 g capped plastic cups) on pallets containing dates of vars. 'Hallawi', 'Hadrawi', 'Zahidi', 'Derei' and 'Ameri'. Dates were sampled before closing the chamber, after one month and after 4.5 months at the end of storage. The following were investigated: presence of insects and molds; skin sloughing; sugar formation; water activity (a_w); and color changes (only on var. Zahidi). At the initial purge phase the desired CO₂ concentration was obtained in the chamber within one hour by introducing the gas under high pressure. An intermittent maintenance phase was then applied for 4.5 months using approximately 0.8 kg CO₂ per day. At the end of storage, quality of the treated dates was compared to controls stored at -18°C. No significant difference was found between the treated dates and controls. The insect population was effectively controlled. This technology is proposed for the treatment of stored dates to control pests and maintain quality.

INTRODUCTION

Dried fruits are subject to infestation by insect pests during and after harvest. Several species of nitidulid beetles are particularly associated with dried fruits because they are both field and storage pests. Fumigation of dried fruits with methyl bromide (MB) upon arrival at the packing plant effectively controls infestation and also causes a high proportion of larvae and adults to emigrate from the fruit before they succumb (Donahaye *et al.* 1991; Navarro *et al.* 1989). The mechanism causing this emigration is not understood, though it clearly involves the delayed action of MB on insects which differs from other fumigants that have an anesthetic effect (Price, 1985).

In recent years, reliance upon fumigation as an overall solution to infestation problems in durable agricultural products has become questionable. The chemical action of fumigants upon commodities and the environment has necessitated the withdrawal of many fumigants from the market. Of the two still widely used, phosphine is characterized as a slow acting fumigant with the ability of insects to develop resistance to it (Winks 1987), whereas MB is under severe scrutiny because

of its involvement in ozone depletion and as such the international community has decided to phase out its use (UNEP, 1998).

The influence of different controlled atmospheres (CAs) in causing emigration of *Carpophilus* spp. larvae from dates was compared with that of MB by Navarro *et al.* (1989) and Donahaye *et al.* (1991). The influence of low O₂ or high CO₂ atmospheres as alternatives to fumigation of dried fruits has been investigated (Soderstrom, 1984; Soderstrom *et al.* 1986; Tarr *et al.* 1994). Recommended dosages to obtain mortality of most stored-product pests using high carbon dioxide (CO₂) CAs require exposure to >60% CO₂ for at least 11 days (Navarro *et al.* 1990).

Generally, soft-fruit date cultivars contain mainly glucose and fructose and little or no sucrose, whereas date cultivars with a firm and dry texture contain a high percentage of sucrose. The sugar that accumulates in the unripe fruits is sucrose and only at a later stage is it inverted. Kanner *et al.* (1978) studied the activity of the invertase enzyme during the last stages of date maturation and found that the maximum invertase activity was much higher in soft-date than in dry-date cultivars. They also discussed the role of invertase in causing sucrose inversion in dates. In soft date cultivars, which contain high amounts of reducing sugars in the ripe fruits, non-enzymatic browning induces development of a "caramel"-like taste (Kanner, 1967). Both accumulation of sugar on the date surface and changes in color are factors that affect date quality during storage (Rygg, 1957).

In Israel, the conventional method of date conservation after harvest is cold-storage at -18°C. This is the most suitable method for the sensitive soft-fruit cultivars, but it is highly energy consuming. In contrast to studies carried out on CAs for insect disinfestation, only very limited work has been carried out to determine the influence of CAs on date quality. Studies under laboratory conditions (Navarro, 1988) and in field tests at ambient temperatures (Navarro *et al.* 1992, 1995) showed that CO₂ significantly delayed browning and sugar formation on dates, and extended shelf life. On the basis of these earlier studies aimed at providing a non-toxic alternative to MB fumigation at ambient temperatures and to extending the shelf life of dates, it was decided to study the application of CA technology to date storage on a commercial scale. This study describes the use of a gastight plastic liner especially designed for the CA storage of dates in Israel.

MATERIALS AND METHODS

Description of the gastight flexible structure (cube)

The flexible structure was erected on a shaded asphalt-paved site and was subjected to the ambient temperatures prevailing in the Bet-Shaan valley in Israel. The completed structure resembled a cube and is therefore referred to as a "cube" in this paper (Fig. 1). It was built up *in-situ* by capping a stack of date crates placed on a floor-liner of PVC sheeting, and then welding the sections into place. It was equipped with a ball valve welded to the lower front side for the introduction of CO₂ and a screw-on valve 100 mm diam. welded at one corner at the top for pressure release. The cube measured 4.2 m high by 6 m wide and 6 m long and had a volume

of 151 m³. These dimensions enabled the storage of 72 crates on pallets each measuring 1 m x 1 m x 2.1 m high.




Fig. 1. Cut-away of cube showing crates on pallets inside cube of 151 m³ capacity.

Test for gastightness

The initial tests for gas tightness were carried out before the cube was loaded with dates, but containing 72 empty crates on pallets so as to enable the shape of the structure to be retained. In this manner, the cube was tested without dates three times with improvements in the gastight seal made between each test, and then once again with the dates inside. A suction fan was used to create a negative pressure, thus enabling gastightness to be tested using the pressure decay method (Annis and van Graver, 1991). A "Celesco" transducer was used to monitor the pressure. The tests were carried out by creating an initial negative pressure of 40 to 60 mm water column, after which the ball valve was closed and the rate of rise in pressure was measured.


Quantity of dates stored in the cube

The dates were stored in the cube in bulk (in boxes of 10 kg or crates of 400 kg on pallets), or packed (250 g capped plastic cups) on pallets. To load the dates into the preformed cube the top section was cut and separated in order to enable removal of 30 empty pallets, which were then replaced with 30 pallets containing dates of the following varieties: Hallawi, Hadrawi, Zahidi, Derei and Ameri. Thus the cube was

partially loaded with 30 tonnes of dates and then sealed, while the remaining space in the cube was filled with 42 pallets without dates.

CA treatments - CO₂ gas purge and maintenance

Four trials were carried out on the purge and maintenance of the CO₂ concentration in the cube without dates and one trial was carried out after loading the cube with dates. In this paper only the trial on the cube with dates is reported. During the purge phase the upper valve was opened in order to prevent build-up of gas pressure. Liquid CO₂ was allowed to flow into the cube from the lower valve, which was connected to one of a series of inverted 24 kg capacity CO₂ gas cylinders (each one emptied into the cube within 4 min). The vaporized gas, entering from the bottom, expelled the air inside the cube upwards and out of the opening at the top (Fig. 2 (A)). Preliminary experiments indicated that uniform CO₂ concentration could be obtained within 48 h of termination of purge. These results are not reported in this paper, but they are schematically shown in Fig. 2 (B). During the maintenance phase the gas was metered into the cube to compensate for small losses through the plastic and through any undetected holes.



Control and monitoring of gas concentrations

The CO₂ concentration during the maintenance phase was controlled using a "David Bishop" O₂ analyzer equipped with control relays capable of operating in the presence of high concentrations of CO₂. The analyzer was set to activate a solenoid valve attached to the CO₂ cylinder when the CO₂ concentration within the cube rose above the set point. The rate of CO₂ flow was determined using a pressure regulator set at a flow-rate between 450 and 940 mL/min. In addition to continuous measurement of O₂ concentration, the CO₂ concentration was measured daily with a portable "GOW-MAC" gas analyzer equipped with a TC detector.

Sampling and test of date quality

Dates were sampled before closing the cube, after one month, and after 4.5 months at the end of storage. Tests included: i) presence of insects and molds according to Israel standard 1251 (Dried Fruits -Dates); ii) skin sloughing; iii) sugar formation; iv) water activity (a_w); and v) color changes (only on var. Zahidi).

RESULTS AND DISCUSSION

Degree of gastightness

The results of tests for gastightness of the cube are presented in Fig. 3. After the second test there was a slight improvement in gastightness. But still the half-life pressure decay was 4.5 min. Following a major repair, after the main leak source was detected, two gastightness tests were carried out, one prior to filling the cube with dates, on March 7, 1995 and the other after loading dates, the next day. These tests showed extended half-life pressure decay of 28 and 16.5 minutes, respectively.




Fig. 3. Pressure decay tests (in mm water column) to evaluate the gastightness of the 151-m³ capacity plastic storage enclosure. The vertical bars indicate the half-life time of pressure decay in each test.

Gas concentrations in the cube

Recordings of CO₂ concentrations in the cube without dates were used to compare the degree of CO₂ loss at different gastightness levels as the sealing of the cube was improved after each test. These results are not reported in this paper. Fig. 4 shows the CO₂ concentrations within the cube containing dates over a period of about 4.5 months. The level of CO₂ remained at ~ 85% for the first 10 days and decreased to ~ 60%, while the O₂ level was about 4% at the outset and rose to 7%. These O₂ concentrations were used as set-points for the monitoring system.

Quantity of gas required to create the CO₂ based controlled atmosphere

To create the CO₂ enriched atmosphere 180 kg CO₂ (equivalent to 98.6 m³ of gas at 30°C) were used in the trial with dates. This is equivalent to an initial CO₂ concentration of 92%. Actually, at the outset an average concentration of 87.8% was recorded in the cube. The difference between the calculated (92%) concentration and the measured average (87.8%), could have been caused by the absorption of the gas by the dates, the wooden pallets, and the plastic boxes.




Fig. 4. Changes in carbon dioxide concentration in the 151 m³ capacity plastic storage enclosure containing 30 tonnes of dates stored for 4.5 months.

Balance of the CO₂ losses during storage

To maintain a constant concentration over 4.5 months of storage 4 CO₂ cylinders per day were required, equivalent to a consumption of 0.8 kg CO₂/day (438 L/day). When liner specification for CO₂ permeability, and the calculated surface area of the cube are taken into account, loss of CO₂ through the plastic cover was calculated to be 202 L/day. The difference between the theoretical amount of gas to be supplied (202 L CO₂/day), and the actual amount required to maintain the target concentration

(438 L/day) indicates the existence of leaks that were not discovered and through which the gas escaped.

The relation between the degree of gastightness and CO₂ losses

To express gastightness in terms of half-life time of pressure decay, the average CO₂ losses per day were calculated after each purge. The results given in Fig. 5, are values for half-life pressure decay plot against the daily % CO₂ losses. It should be pointed out that this test is dependent on the free volume of the structure. From the shape of the curve, it is possible to conclude that in structures of similar size, a half-life pressure decay of less than 10 min can be regarded as critical, since less gastight structures would require the use of large amounts of CO₂ to maintain a constant concentration.




Fig. 5. Degree of gastightness expressed in half-life (min) pressure decay and daily percentage carbon dioxide loss from the 151-m³ capacity storage enclosure.

CA storage for quality preservation of dates

In this trial date quality was preserved under CA at temperatures ranging between 22 and 28°C. Tests for presence of insects and molds revealed no development during storage in the cube. Water activity of dates remained low from beginning to end of storage in the range of 0.55 to 0.64 *a_w*.

No significant change in sloughing of peel, or in sugar formation on the date surface was observed after 4.5 months of storage. These results support data obtained in a previous study of Zahidi, Ameri and Hallawi dates stored under a CO₂ enriched CA, namely that there was no negative influence as regards increase of sugars, even after 8 months of storage (Navarro *et al.* 1992; 1995).

CONCLUSIONS

Our results showed: i) that the use of CA storage without cooling is feasible; ii) it was found necessary to supply 0.8 kg/day of CO₂ to maintain a concentration of 60 to 80% in the cube; iii) to prevent further CO₂ loss the degree of gas tightness should ensure a half-life pressure decay of not less than 10 min; iv) the level of moisture in the dates did not change and no reduction in quality was observed throughout the 4.5 month storage period.

ACKNOWLEDGEMENTS

The authors thank the chief technologist of Tamar Shaan Eng. Dov Avni and Mrs Judy Zeira for their assistance during the monitoring tests of the cube, as well as the manager of Tamar Shaan Mr. Itshak Sagi and the staff of the packing house for their support throughout the trials. Likewise we thank Haogenplast Ltd. for supplying the plastic cube and especially Mr. Tom deBruin, marketing manager of Haogenplast. This project (no. 417-0214-96) was supported by the Chief Scientist of the Israel Ministry of Agriculture and, the R&D Energy Conservation Committee in Agriculture.

REFERENCES

- Annis, P.C. and van S. Graver, J. (1991) Suggested recommendations for the fumigation of grain in the ASEAN region. Part 2. Carbon dioxide fumigation of bag-stacks sealed in plastic enclosures: an operations manual. ASEAN Food Handling Bureau Kuala Lumpur, Malaysia and ACIAR, Canberra, Australia 58 pp.
- Donahaye, J.E., Navarro, S., Rindner, M. and Dias, R. (1991) The influence of different treatments causing emigration of nitidulid beetles. *Phytoparasitica* **19**, 273-282.
- Israel Standard No. 1251, (1984) Dates. Israel Institute of Standards, 7 pp.
- Kanner, J. (1967) Rehydration of overdried dates and biochemical changes involved in the rehydrated fruits. M. Sc. Thesis, Faculty of Agriculture of the Hebrew University of Jerusalem (in Hebrew with English summary).
- Kanner J., Elmaleh, H., Reuveni, O. and Ben-Gera, I. (1978) Invertase (β -fructofuranosidase) activity in three date cultivars. *J. Agric. Food Chem.*, **26**, 1239-1240.
- Navarro, S. and Donahaye, E. (1990) Generation and application of modified atmospheres and fumigants for the control of storage insects. In: *Fumigation and Controlled Atmosphere Storage of Grain: Proc. Int. Conf.* (Edited by Champ, B.R., Highley, E., and Banks, H.J.) Singapore, 14-18 Feb. 1989 ACIAR Proceedings No. 25. 152-165.
- Navarro, S., Donahaye, E., Azrieli, A., Rindner, M. and Silberstein, B. (1992) Energy saving by storing dates without cooling. Ann. Rep. Project No. 417-0214-91. Israel Agric. Res. Organ., Bet-Dagan, 11 pp (in Hebrew).

- Navarro, S., Donahaye, E., Silberstein, B., Azrieli, A. and Rindner M. (1995) Energy saving by storing dates without cooling. Ann. Rep. Project No. 417-0214-94. Israel Agric. Res. Organ., Bet-Dagan, 21 pp (in Hebrew).
- Navarro, S., Paran, G., Azrieli, A. and Donahaye, E. (1988) Influence of carbon dioxide on color preservation of deglet noor dates. Ann. Rep. Project No. 564-001-87. Israel Agric. Res. Organ., Bet-Dagan, 10 pp (in Hebrew).
- Navarro, S., Donahaye, E., Dias, and Jay, E. (1989) Integration of modified atmospheres for disinfestation of dried fruits. Final Rep. of BARD Project No. I-1095-86, 86 pp.
- Price, N. R. (1985) The mode of action of fumigants. *J. stored Prod. Res.*, **21**, 157-164.
- Rygg, G.L. (1957) The relation of moisture content to rate of darkening in deglet noor dates. *Date Growers Inst. Ann. Rept.*, **34**, 13-14.
- Soderstrom, E.L. (1984) Economic cost evaluation of a generated low-oxygen atmosphere as an alternative fumigant in the bulk storage of raisins. *J. Econ Entomol.*, **77**, 457-461.
- Soderstrom, E.L., Mackey, B.E. and Brandl, D.G. (1986) Interactive effects of low-oxygen atmospheres, relative humidity and temperature on mortality of two stored product moths (Lepidoptera, Pyralidae). *J. Econ Entomol.*, **79**, 1303-1306.
- Tarr, C., Hilton, S.J., van S. Graver J. and Clingeffer P.R. (1994) Carbon dioxide fumigation of processed dried vine fruit (sultanas) in sealed stacks. In: *Proc. 6th Int. Working Conf. on Stored-Product Protection* (Edited by Highley, E., Wright, E.J., Banks, H.J. and Champ, B.R.) Canberra, Australia, 17-23 April 1994, CAB International, Wallingford, Oxon, UK, **1**, 204-209.
- UNEP (1998). Montreal Protocol on Substances that deplete the Ozone Layer. 1998 Assessment of Alternatives to Methyl Bromide. Methyl Bromide Alternatives Committee. Nairobi, Kenya.
- Winks, R.G. (1987) Strategies for the effective use of phosphine as a grain fumigant and the implications of resistance. In: *Proc. 4th Int. Working Conf. on Stored-product Protection*, (Edited by Donahaye, E., Navarro, S.), Tel Aviv. Caspit Press, Jerusalem, 335-344.